


*Europ. J. Combinatorics* (2001) **22**, 415–429

doi:10.1006/eujc.2000.0466

Available online at <http://www.idealibrary.com> on  IDEAL<sup>®</sup>

## Exact $m$ -covers of Groups by Cosets

ZHI-WEI SUN

Let  $G$  be a group covered by its left cosets  $a_1 G_1, \dots, a_k G_k$  exactly  $m$  times. It is known that  $[G : \bigcap_{i=1}^k G_i] \leq k!$ . When all the  $G_i$  are subnormal in  $G$  and  $\bigcap_{i=1}^k G_i = H$ , we are able to determine the least value of  $k$  in terms of  $m, G, H$ . For any  $i = 1, \dots, k$ , providing  $G/(G_i)_G$  is solvable we show that  $k \geq m + f([G : G_i])$  and hence  $[G : G_i] \leq 2^{k-m}$ , where  $f(n) = \sum_{s=1}^r \alpha_s(p_s - 1)$  if  $p_1^{\alpha_1} \cdots p_r^{\alpha_r}$  is the standard factorization of  $n$ . These extend some previous results on disjoint covers.

© 2001 Academic Press

### 1. INTRODUCTION

Let  $G$  be a (multiplicative) group (whose identity element is denoted by  $e$ ) and  $H$  a subgroup of  $G$ . For any  $a \in G$  we call  $aH = \{ax : x \in H\}$  a left coset of  $H$  in  $G$ , by  $G/H$  we mean the set of all left cosets of  $H$  in  $G$ . As usual, the index of  $H$  in  $G$  is  $[G : H] = |G/H|$ , and  $G/H$  is called the quotient group (or the factor group) of  $G$  by  $H$  if  $H$  is normal in  $G$ . We use  $H_G$  and  $H^G$  to denote the core (i.e., normal interior) and the normal closure of  $H$  in  $G$ , respectively.

Let  $a_1 G_1, \dots, a_k G_k$  be finitely many left cosets in a group  $G$ . For the system

$$\mathcal{A} = \{a_i G_i\}_{i=1}^k, \quad (1.1)$$

we call

$$w_{\mathcal{A}}(x) = |\{1 \leq i \leq k : x \in a_i G_i\}| \quad (1.2)$$

the covering multiplicity of  $x \in G$ , and put  $m(\mathcal{A}) = \inf_{x \in G} w_{\mathcal{A}}(x)$ .

Let  $m \in \mathbb{Z}^+ = \{1, 2, 3, \dots\}$ . (1.1) is said to be an exact  $m$ -cover of  $G$  if  $w_{\mathcal{A}}(x) = m$  for all  $x \in G$ . Exact 1-covers are just partitions into left cosets, they are also called *disjoint covers*. As mentioned in all textbooks on group theory, for any subgroup  $H$  of  $G$  with finite index, all the  $[G : H]$  left cosets of  $H$  in  $G$  form a disjoint cover of  $G$ . Clearly the only exact  $m$ -cover of  $G$  by subgroups consists of  $m$  copies of  $G$ . It is known that even for the additive group  $\mathbb{Z}$  of integers there exists an exact  $m$ -cover no subsystem of which forms an exact  $n$ -cover with  $0 < n < m$  (cf. [4, 24]). Exact  $m$ -covers are natural extension of disjoint covers, they should have regular properties because any such a cover covers all the elements the same times. Covers of  $\mathbb{Z}$  by residue classes were initiated by Erdős [2] (see Section 1 of Sun [20] for problems and results in the area), exact  $m$ -covers of  $\mathbb{Z}$  were first investigated by Porubský [14], exact  $m$ -covers of an arbitrary group have been studied only in the case  $m = 1$ .

Exact 1-covers are very special in the sense that members of such a cover are pairwise disjoint. There has been much research on such covers, see [12, 15, 27] for results concerning disjoint covers of  $\mathbb{Z}$ . Often there are no obvious ways to generalize results on disjoint covers to exact  $m$ -covers, this is why few properties of exact  $m$ -covers are known. By an easy counting argument, if a system

$$\mathcal{A} = \{a_i + n_i \mathbb{Z}\}_{i=1}^k \quad (a_1, \dots, a_k \in \mathbb{Z}; \quad n_1, \dots, n_k \in \mathbb{Z}^+) \quad (1.3)$$

of residue classes forms an exact  $m$ -cover of  $\mathbb{Z}$  then  $\sum_{i=1}^k 1/n_i = m$ . In [19–21] the author revealed some further connections between exact  $m$ -covers of  $\mathbb{Z}$  and unit fractions. In this paper we shall prove some inequalities for exact  $m$ -covers of groups.

The Mycielski function  $f : \mathbb{Z}^+ \rightarrow 0, 1, 2, \dots$  is determined as follows:

$$f(p) = p - 1 \text{ for any prime } p \text{ and } f(mn) = f(m) + f(n) \text{ for all } m, n \in \mathbb{Z}^+.$$

Evidently

$$f(p_1^{\alpha_1} \cdots p_r^{\alpha_r}) = \sum_{s=1}^r \alpha_s (p_s - 1) \quad (1.4)$$

where  $p_1, \dots, p_r$  are distinct primes and  $\alpha_1, \dots, \alpha_r$  are non-negative integers. In 1966 Mycielski and Sierpiński [9] conjectured that if (1.1) forms a disjoint cover of an abelian group  $G$  and all the  $[G : G_i]$  are finite then

$$k \geq 1 + \max_{1 \leq i \leq k} f([G : G_i]). \quad (1.5)$$

This was confirmed by Znám [25] for  $G = \mathbb{Z}$ , and verified by Hejny and Znám [5] in a special case. In 1968 Znám [26] posed a conjecture that if (1.3) forms a disjoint cover of  $\mathbb{Z}$  then  $k \geq 1 + f(N)$  where  $N$  is the least common multiple of  $n_1, \dots, n_k$ . Later in 1974 his student Korec ([6]) obtained the following stronger result: Let  $G$  be a group and (1.1) its disjoint cover with all the  $G_i$  normal in  $G$ , then each  $G_i$  has finite index in  $G$  and

$$k \geq 1 + f\left(\left[G : \bigcap_{i=1}^k G_i\right]\right). \quad (1.6)$$

In 1988 Berger *et al.* [1] proved that (1.5) holds if (1.1) forms a disjoint cover of a finite solvable group  $G$ .

It seems that Mycielski, Znám and Korec had not known the following basic result established by Neumann [10, 11] in 1954: if (1.1) forms a cover of a group  $G$  then  $G$  is the union of those  $a_i G_i$  with  $[G : G_i] < \infty$ . In 1987 Tomkinson [23] strengthened the Neumann result by showing that if (1.1) forms a cover of a group  $G$  but none of its proper subsystems does then

$$\left[G : \bigcap_{i=1}^k G_i\right] \leq k! \quad (1.7)$$

where the upper bound  $k!$  is the best possible. By Corollary 1 of Sun [18], (1.7) holds if  $m(\mathcal{A}') < m(\mathcal{A}) = m$  for any proper subsystem  $\mathcal{A}'$  of  $\mathcal{A}$ . In particular, when (1.1) forms an exact  $m$ -cover of group  $G$ , all the  $G_i$  and hence the intersection  $\bigcap_{i=1}^k G_i$  are of finite index in  $G$ .

Let  $G$  be a group and  $H$  a subnormal subgroup of  $G$  with finite index. Let

$$H_0 = H \subset H_1 \subset \cdots \subset H_n = G$$

be a composition series from  $H$  to  $G$ . If the length  $n$  is zero (i.e.,  $H = G$ ), then we set  $d(G, H) = 0$ , otherwise we put

$$d(G, H) = \sum_{i=0}^{n-1} ([H_{i+1} : H_i] - 1). \quad (1.8)$$

By the Jordan–Hölder theorem,  $d(G, H)$  does not depend on the choice of the composition series from  $H$  to  $G$ . Theorem 6 of [18] indicates that

$$[G : H] - 1 \geq d(G, H) \geq f([G : H]) \geq \log_2 [G : H]. \quad (1.9)$$

In contrast with Tomkinson's result, inequality (1.6) implies that

$$\left[ G : \bigcap_{i=1}^k G_i \right] \leq 2^{k-1}. \quad (1.10)$$

In 1990 the author [18] obtained the following improvement to Korec's result: if (1.1) forms a disjoint cover of a group  $G$  with all the  $G_i$  subnormal in  $G$ , then

$$k \geq 1 + d\left(G, \bigcap_{i=1}^k G_i\right). \quad (1.11)$$

Let us look at two examples.

EXAMPLE 1.1. Let  $G$  be a group and  $H$  a subgroup with  $k = [G : H] < \infty$ . Suppose that  $\{Ha_1, \dots, Ha_k\}$  is a right coset decomposition of  $G$ . Set  $G_i = a_i^{-1}Ha_i$  for  $i = 1, \dots, k$ . The author [18] observed that (1.1) forms a disjoint cover of  $G$  with  $\bigcap_{i=1}^k G_i = H_G$ . When  $G = S_k$  is the symmetric group on  $\{1, \dots, k\}$  and  $H$  is the stabilizer of 1,

$$\{H, H(12), H(13), \dots, H(1k)\} = \{G_1, (12)G_2, \dots, (1k)G_k\}$$

forms a partition of  $G$  where  $G_i$  is the stabilizer of  $i$  for each  $i = 1, \dots, k$ , Tomkinson [23] noticed that in this case  $\bigcap_{i=1}^k G_i = H_G = \{e\}$  has index  $k!$  in  $G$ ; we add here that if  $k \geq 3$  then all the subgroups  $G_1, \dots, G_k$  are distinct,  $[G : \bigcap_{i=1}^k G_i] = k! (> 2^{k-1})$  has some prime divisors (e.g., prime factors of  $k-1$ ) not dividing  $k = [G : G_i]$ , and  $f([G : \bigcap_{i=1}^k G_i]) = \sum_{i=2}^k f(i) \geq f(3) + k - 2 = k$ .

EXAMPLE 1.2. Let  $G$  be a group and  $H = H_0 \subset H_1 \subset \dots \subset H_n = G$  be a chain of subgroups with finite index. Let  $a \in G$  and  $H_{i+1} \setminus H_i = \bigcup_{j=1}^{[H_{i+1}:H_i]-1} b_j^{(i)} H_i$  for  $i = 0, 1, \dots, n-1$ . In [18] the author observed that those  $ab_j^{(i)} H_i$  ( $0 \leq i < n$  and  $1 \leq j < [H_{i+1} : H_i]$ ) form a partition of  $G \setminus aH_0$ . Thus, the cosets, together with  $aH_0$  and  $m-1$  copies of  $G$ , form an exact  $m$ -cover of  $G$  with the number  $k$  of cosets being  $m + \sum_{i=0}^{n-1} ([H_{i+1} : H_i] - 1)$  and the intersection of the  $k$  subgroups being  $H$ . If  $H_i$  is maximal normal in  $H_{i+1}$  for each  $i = 0, \dots, n-1$  then  $k = m + d(G, H)$ . In general, as  $[H_{i+1} : H_i] - 1 \geq f([H_{i+1} : H_i])$  we have

$$k - m \geq \sum_{i=0}^{n-1} f([H_{i+1} : H_i]) = f\left(\prod_{i=0}^{n-1} [H_{i+1} : H_i]\right) = f([G : H]).$$

In this paper we study lower bounds for the number  $k$  of cosets in an exact  $m$ -cover (1.1) of a group  $G$ , with the intersection  $\bigcap_{i=1}^k G_i$  or a subgroup  $G_i$  given. In the next section we prove a key property of exact  $m$ -covers. In Section 3 we characterize those subnormal subgroups  $H$  of group  $G$  for which  $[G : H] < \infty$  and  $d(G, H) = f([G : H])$ . In Section 4 we present our main results with an application in group theory.

Now we state our central results. (Actually we will prove more.)

(I) Let  $G$  be a group and (1.1) an exact  $m$ -cover of  $G$  with all the  $G_i$  subnormal in  $G$ . Then  $k \geq m + d(G, \bigcap_{i=1}^k G_i)$ . Moreover, for any subgroup  $K$  of  $G$  not contained in all the  $G_i$  we have

$$|\{1 \leq i \leq k : K \not\subseteq G_i\}| \geq 1 + d\left(K, K \cap \bigcap_{i=1}^k G_i\right).$$

(II) Let (1.1) be an exact  $m$ -cover of a group  $G$ . Whenever  $G/(G_i)_G$  is solvable, we have  $k \geq m + f([G : G_i])$  and hence  $[G : G_i] \leq 2^{k-m}$ .

Concerning result (II) we have a further conjecture.

CONJECTURE . Let (1.1) be an exact  $m$ -cover of a group  $G$  with all the  $G/(G_i)_G$  solvable. Then  $k \geq m + f(N)$  where  $N$  is the least common multiple of  $[G : G_1], \dots, [G : G_k]$ .

For a cover (1.1) of group  $G$ , if it does not form an exact  $m$ -cover for any  $m = 1, 2, 3, \dots$ , then we do not have a similar inequality in general. When  $G$  is cyclic, or  $|G|$  is square free and all the  $G_i$  are subnormal in  $G$ , if  $m(\mathcal{A}') < m(\mathcal{A})$  for any proper subsystem  $\mathcal{A}'$  of  $\mathcal{A}$  then we can show that  $k \geq m(\mathcal{A}) + f([G : \bigcap_{i=1}^k G_i])$ . Since this result and its extension are quite involved, we do not present a proof here.

## 2. A VITAL PROPERTY OF EXACT $m$ -COVERS

Of course, whether two left cosets of subgroups are disjoint, is very essential. Here we give the following lemma.

LEMMA 2.1. Let  $H$  and  $K$  be subgroups of a group  $G$ . Then:

- (i)  $HK = G$  if and only if  $xH \cap yK \neq \emptyset$  for all  $x, y \in G$ .
- (ii)  $HK = KH$  coincides with  $G$  or  $H$  in the following cases:
  - (a)  $H$  is maximal and normal in  $G$ ;
  - (b)  $H$  is maximal in  $G$  and  $K$  is normal in  $G$ ;
  - (c)  $H$  is maximal normal and  $K$  is subnormal in  $G$ .

PROOF. (i) If  $HK = G$ , then for any  $x, y \in G$  there exist  $h \in H$  and  $k \in K$  such that  $hk = x^{-1}y$  and hence

$$xH \cap yK = xhH \cap xhkK = xh(H \cap K) \neq \emptyset.$$

Conversely, if  $xH \cap yK \neq \emptyset$  for all  $x, y \in G$ , then for any  $g \in G$  there is a  $h \in H$  for which  $h \in gK$  and hence  $gK = hK$ , therefore  $G = HK$ .

(ii) In each case  $H$  or  $K$  is normal in  $G$  and thus  $HK = KH$  is a subgroup of  $G$  containing  $H$ . Due to the maximality of  $H$ ,  $HK$  coincides with  $G$  or  $H$  in the cases (a) and (b). In case (c),  $HK$  is subnormal in  $G$  (by 7.19 of [17]), if  $HK \neq G$  then  $(HK)^G \neq G$  and so  $(HK)^G = H$  (i.e.,  $HK = H$ ) by the maximal normality of  $H$ .  $\square$

REMARK 2.1. For previous combined use of parts (i) and (ii), the reader may consult the proofs of Lemma 6 of [6], Theorem 1 of [13], Lemma 2.II of [1] and Theorem 4 of [18]. For a proper subgroup  $H$  of group  $G$ , case (a) is equivalent to the following:  $H$  is a (maximal) normal subgroup of prime index in  $G$ . (Notice that  $H$  is maximal in  $G$  if it has prime index in  $G$ , and that in case (a)  $G/H$  is a cyclic group of prime order since  $G/H$  and  $H/H$  are the only subgroups of  $G/H$ .)

LEMMA 2.2. Let  $G$  be a group and  $G_1, \dots, G_k$  be subgroups of  $G$  with finite index. Then

$$\sum_{i=1}^k \frac{1}{[G : G_i]} = \frac{1}{[G : \bigcap_{j=1}^k G_j]} \sum_{C \in G / \bigcap_{j=1}^k G_j} |\{1 \leq i \leq k : C \subseteq a_i G_i\}| \geq m(\mathcal{A}). \quad (2.1)$$

Therefore, (1.1) forms an exact  $m$ -cover of  $G$  if and only if

$$\sum_{i=1}^k \frac{1}{[G : G_i]} = m \quad (2.2)$$

and  $w_{\mathcal{A}}(x) \leq m$  (or  $w_{\mathcal{A}}(x) \geq m$ ) for all  $x \in G$ .

PROOF.  $H = \bigcap_{j=1}^k G_j$  is of finite index in  $G$ . Clearly

$$[G : H] \sum_{i=1}^k \frac{1}{[G : G_i]} = \sum_{i=1}^k [G_i : H] = \sum_{i=1}^k \sum_{\substack{C \in G/H \\ C \subseteq a_i G_i}} 1 = \sum_{C \in G/H} \sum_{\substack{i=1 \\ a_i G_i \supseteq C}}^k 1.$$

So

$$\begin{aligned} \sum_{i=1}^k \frac{1}{[G : G_i]} &= \frac{1}{[G : H]} \sum_{C \in G/H} |\{1 \leq i \leq k : C \subseteq a_i G_i\}| \\ &\geq \frac{1}{[G : H]} \sum_{C \in G/H} m(\mathcal{A}) = m(\mathcal{A}). \end{aligned}$$

If (1.1) forms an exact  $m$ -cover of  $G$ , then  $|\{1 \leq i \leq k : C \subseteq a_i G_i\}| = m$  for each  $C \in G/H$  and hence (2.2) holds. When  $w_{\mathcal{A}}(x) \leq m$  (or  $w_{\mathcal{A}}(x) \geq m$ ) for all  $x \in G$ , if (2.2) is valid then  $|\{1 \leq i \leq k : C \subseteq a_i G_i\}|$  must coincide with  $m$  for every  $C \in G/H$  and hence  $w_{\mathcal{A}}(x) = m$  for any  $x \in G$ . This completes the proof.  $\square$

REMARK 2.2. It has been known that if (1.3) is an exact  $m$ -cover of  $\mathbb{Z}$  then  $\sum_{i=1}^k 1/[\mathbb{Z} : n_i \mathbb{Z}]$  equals  $m$  (see [14]). In 1977 Korec and Znám [8] proved that  $\sum_{i=1}^k 1/[G : G_i] = 1$  for any disjoint cover (1.1) of group  $G$ .

THEOREM 2.1. Assume that (1.1) forms an exact  $m$ -cover of a group  $G$  (by left cosets of subgroups  $G_1, \dots, G_k$ ). For a subgroup  $H$  of  $G$  we have

$$\{C \in G/H : C \supseteq a_i G_i \text{ for some } i = 1, \dots, k\} = \emptyset \quad \text{or} \quad G/H. \quad (2.3)$$

in the following cases:

- (a)  $H$  is the group  $G$  or a normal subgroup of prime index in  $G$ ;
- (b)  $G_1, \dots, G_k$  are normal in  $G$  and  $H$  is maximal in  $G$ ;
- (c)  $G_1, \dots, G_k$  are subnormal and  $H$  is maximal normal in  $G$ .

PROOF. Suppose, in contrast, that (2.3) is false. Then there exist  $a, b \in G$  such that  $a_j G_j \subseteq aH$  for some  $1 \leq j \leq k$  and that  $a_i G_i \not\subseteq bH$  for all  $i = 1, \dots, k$ . Let  $I = \{1 \leq i \leq k : G_i \not\subseteq H\}$ . Clearly,  $j \notin I$  since  $a_j H = aH \supseteq a_j G_j$ . When  $i \in I$ , by Lemma 2.1 and Remark 2.1 we have  $G_i H = G$  and  $a_i G_i \cap bH \neq \emptyset$  in all the three cases. If  $s \in \{1, \dots, k\} \setminus I$ , then  $a_s G_s \subseteq a_s H \neq bH$  and hence  $a_s G_s \cap bH = \emptyset$ . As (1.1) forms an exact  $m$ -cover of  $G$ ,  $I$  must be non-empty and  $\{b^{-1}a_i G_i \cap H\}_{i \in I}$  must be an exact  $m$ -cover of  $H$  by left cosets of subgroups  $G_i \cap H$  ( $i \in I$ ) in  $H$ . In view of Lemma 2.2,

$$\sum_{i \in I} \frac{1}{[H : G_i \cap H]} = m = \sum_{i=1}^k \frac{1}{[G : G_i]}.$$

However, for each  $i \in I$  we have  $[H : G_i \cap H] = [G : G_i]$  because  $G = G_i H$  contains exactly  $[H : G_i \cap H]$  right cosets of  $G_i$  by Chapter 1 of [22]. So  $I$  must coincide with  $\{1, \dots, k\}$ , which contradicts the fact that  $j \notin I$ . The proof is complete.  $\square$

REMARK 2.3. When  $m = 1$ , Theorem 2.1 in case (c) was obtained by the author [18] in a particular way.

3. WHEN  $d(G, H) = f([G : H])$ ?

For any group  $G$  we let  $\mathcal{S}(G)$  denote the class of subnormal subgroups  $H$  of  $G$  for which  $[G : H] < \infty$  and  $d(G, H) = f([G : H])$ . In this section we aim to characterize those  $H \in \mathcal{S}(G)$ .

LEMMA 3.1. *Let  $G$  be a group.*

- (i) *For subgroups  $H$  and  $K$  of  $G$  with  $[G : H]$  finite and  $H$  or  $K$  subnormal in  $G$ , we have  $[K : H \cap K] \mid [G : H]$ .*
- (ii) *If  $G_1, \dots, G_k$  are subnormal subgroups of  $G$  with finite index, then*

$$\left[ G : \bigcap_{i=1}^k G_i \right] \mid \prod_{i=1}^k [G : G_i]. \quad (3.1)$$

PROOF. (i) When  $H$  is subnormal in  $G$ , there exists a finite chain

$$H = H_0 \subseteq H_1 \subseteq \dots \subseteq H_{n-1} \subseteq H_n = G$$

of subgroups of  $G$  such that  $H_{i-1}$  is normal in  $H_i$  for all  $i = 1, \dots, n$ . By the second isomorphism theorem (for group  $H_i$ ) (cf. 3.40 of [17])

$$[H_i \cap K : (H_i \cap K) \cap H_{i-1}] = |(H_i \cap K)H_{i-1}/H_{i-1}|$$

and hence  $[H_i \cap K : H_{i-1} \cap K] \mid [H_i : H_{i-1}]$ . We therefore have

$$\prod_{i=1}^n [H_i \cap K : H_{i-1} \cap K] \mid \prod_{i=1}^n [H_i : H_{i-1}], \text{ i.e., } [K : H \cap K] \mid [G : H].$$

Now we assume that  $K$  is subnormal in  $G$ . Let  $K_0 = K \subseteq K_1 \subseteq \dots \subseteq K_n = G$  be a chain of subgroups such that  $K_{i-1}$  is normal in  $K_i$  for any  $i = 1, \dots, n$ . In view of Chapter 1 of [22], the subgroup  $K_{i-1}(H \cap K_i)$  of  $K_i$  contains exactly  $[K_{i-1} : K_{i-1} \cap (H \cap K_i)] = [K_{i-1} : H \cap K_{i-1}]$  left cosets of  $H \cap K_i$  and so  $[K_{i-1} : H \cap K_{i-1}] \mid [K_i : H \cap K_i]$ . It follows that  $[K : H \cap K] = [K_0 : H \cap K_0]$  divides  $[K_n : H \cap K_n] = [G : H]$ .

This proves the first part.

(ii) By part (i), if  $H$  and  $K$  are subnormal subgroups of  $G$  with finite index, then  $[G : H \cap K] = [G : K][K : H \cap K]$  divides  $[G : H][G : K]$ . Thus we can easily show part (ii) by induction. This completes the proof.  $\square$

REMARK 3.1. Let  $G$  be a group and  $H, K$  be subgroups of  $G$  with  $[G : H] < \infty$ . Although  $[K : H \cap K] \leq [G : H]$ , in general we may have  $[K : H \cap K] \nmid [G : H]$  even if  $G$  is solvable. For example, the symmetric group  $G = S_3$  on  $\{1, 2, 3\}$  has subgroups  $H = \{e, (12)\}$  and  $K = \{e, (13)\}$  with  $H \cap K = \{e\}$ , apparently  $[K : H \cap K] = |K| = 2$ ,  $[G : H] = 3!/2 = 3$  and  $[G : H \cap K] \nmid [G : H][G : K]$ .

Recall that a group  $G$  is called *perfect* if  $G$  coincides with its commutator subgroup  $G' = [G, G]$ , and that every finite group has a unique solvable residual (see 7.50 of [17]).

THEOREM 3.1. *Let  $G$  be a group and  $H$  be a subgroup of  $G$ .*

- (i)  *$H \in \mathcal{S}(G)$  if and only if there is a composition series from  $H$  to  $G$  whose factors are of prime orders.*
- (ii) *When  $H \in \mathcal{S}(G)$ , we have  $H \cap K \in \mathcal{S}(K)$  for any subgroup  $K$  of  $G$ , also  $H \cap K, \langle H, K \rangle \in \mathcal{S}(G)$  if  $K \in \mathcal{S}(G)$ .*

- (iii) If  $H$  lies in  $\mathcal{S}(G)$  then so do  $H_G$  and  $H^G$ .
- (iv)  $H \in \mathcal{S}(G)$  if and only if  $H$  is subnormal in  $G$  and  $G/H_G$  is finite and solvable.
- (v) If  $H \in \mathcal{S}(G)$  then  $H$  contains all perfect subgroups of  $G$ . When  $G$  is finite and  $H$  is subnormal in  $G$ , we have  $H \in \mathcal{S}(G)$  if  $H$  contains the (perfect) normal subgroup  $K$  of  $G$  for which  $G/K$  is the solvable residual of  $G$ .
- (vi) Assume that  $H$  is subnormal in  $G$  and  $[G : H]$  is finite. Then  $H \in \mathcal{S}(G)$  if  $G$  is locally solvable, or  $[G : H]$  is square free or odd or divisible by at most two distinct primes.

PROOF. (i) In the case  $H = G$ , clearly  $H \in \mathcal{S}(G)$  since  $d(G, H) = 0 = f([G : H])$ , and the composition series from  $H$  to  $G$  has length zero and no factor.

Now let  $H$  be proper in  $G$ . Apparently  $H$  is subnormal and of finite index in  $G$  if and only if there is a composition series from  $H$  to  $G$  whose factors are finite. Suppose that  $H$  is such a subgroup of  $G$  and that  $H = H_0 \subset H_1 \subset \cdots \subset H_n = G$  is a composition series from  $H$  to  $G$ . Then (1.8) holds and

$$f([G : H]) = f\left(\prod_{i=1}^n |H_i/H_{i-1}|\right) = \sum_{i=1}^n f(|H_i/H_{i-1}|).$$

Since  $m - 1 \geq f(m)$  for all  $m \in \mathbb{Z}^+$ , and

$$f(m_1 m_2) = f(m_1) + f(m_2) \leq m_1 - 1 + m_2 - 1 < m_1 m_2 - 1$$

for any integers  $m_1, m_2 > 1$ , we therefore have

$$\begin{aligned} d(G, H) = f([G : H]) &\iff |H_i/H_{i-1}| - 1 = f(|H_i/H_{i-1}|) \text{ for all } i = 1, \dots, n \\ &\iff |H_i/H_{i-1}| \text{ is a prime number for any } i = 1, \dots, n. \end{aligned}$$

This proves part (i).

(ii) When  $H = G$ , part (ii) is obvious. Assume that  $H \in \mathcal{S}(G)$  but  $H \neq G$ . By part (i) there exists a composition series  $H = H_0 \subset H_1 \subset \cdots \subset H_n = G$  from  $H$  to  $G$  such that  $|H_1/H_0|, \dots, |H_n/H_{n-1}|$  are primes. Let  $K$  be a subgroup of  $G$ . For each  $i = 1, \dots, n$ , evidently  $H_{i-1} \cap K = H_{i-1} \cap (H_i \cap K)$  is normal in  $H_i \cap K$  since  $H_{i-1}$  is normal in  $H_i$ ; by Lemma 3.1(i)  $[H_i \cap K : H_{i-1} \cap K]$  divides  $[H_i : H_{i-1}]$  and hence coincides with one or the prime  $|H_i/H_{i-1}|$ . Thus, in view of part (i),  $H \cap K = H_0 \cap K \in \mathcal{S}(H_n \cap K) = \mathcal{S}(K)$ .

Now suppose that  $K \in \mathcal{S}(G)$ . Then  $H \cap K$  is subnormal and of finite index in  $G$ . As  $H \cap K \in \mathcal{S}(K)$  and  $K \in \mathcal{S}(G)$ , we have  $H \cap K \in \mathcal{S}(G)$  by part (i). Put  $L = (H \cap K)_G$ . Then  $H/L, K/L$  and their join  $\langle H/L, K/L \rangle = \langle H, K \rangle/L$  are subnormal in the finite group  $G/L$  (cf. 1.8 and 7.22 of [17]), so  $\langle H, K \rangle$  is subnormal in  $G$ . Since  $(H \cap K)/L \in \mathcal{S}(K/L)$ , by part (i) and 7.25 of [17] we have  $H/L \in \mathcal{S}(\langle H, K \rangle/L)$  and hence  $H \in \mathcal{S}(\langle H, K \rangle)$ . As  $H \in \mathcal{S}(G)$ ,  $\langle H, K \rangle$  must lie in  $\mathcal{S}(G)$ .

(iii) Let  $H \in \mathcal{S}(G)$ . By part (i)  $g^{-1}Hg \in \mathcal{S}(G)$  for any  $g \in G$ . As  $[G : H] < \infty$  there are only finitely many distinct conjugates of  $H$ . Their intersection is  $H_G$  and their join is  $H^G$ . Applying part (ii) we obtain that  $H_G, H^G \in \mathcal{S}(G)$ .

(iv) Let  $H$  be subnormal in  $G$ . By part (iii),  $H_G \in \mathcal{S}(G)$  providing  $H \in \mathcal{S}(G)$ . By part (i), if  $H_G \in \mathcal{S}(G)$  then  $H \in \mathcal{S}(G)$  and  $H_G \in \mathcal{S}(H)$ . So  $H \in \mathcal{S}(G)$  if and only if  $H_G \in \mathcal{S}(G)$ .

Note that  $G/H_G$  is a finite solvable group if and only if there is a composition series from  $H_G/H_G$  to  $G/H_G$  whose factors are of prime orders (cf. 7.56 of [17]). Thus, by part (i),  $H_G \in \mathcal{S}(G)$  (i.e.,  $H \in \mathcal{S}(G)$ ) if and only if  $G/H_G$  is finite and solvable.

(v) As usual, for any group  $F$  we let  $F^{(0)} = F$  and  $F^{(n+1)} = (F^{(n)})'$  for  $n = 0, 1, 2, \dots$ . Suppose  $H \in \mathcal{S}(G)$ . Then  $G/H_G$  is solvable by part (iv), therefore  $(G/H_G)^{(n)} = H_G/H_G$  for

some non-negative integer  $n$  (see 7.52 of [17]). Notice that  $(G/H_G)^{(n)} = (GH_G/H_G)^{(n)} = G^{(n)}H_G/H_G$  (cf. Exercise 164 of [17]). So  $G^{(n)} \subseteq H_G \subseteq H$ . If  $M$  is a perfect subgroup of  $G$ , then  $M = M^{(n)} \subseteq G^{(n)} \subseteq H$ .

For the rest of part (v), let  $G$  be finite, and  $H$  be a subnormal subgroup of  $G$  which contains the smallest normal subgroup  $K$  of  $G$  such that  $G/K$  is solvable. As a characteristic subgroup of  $K$ ,  $K'$  is also normal in  $G$  (cf. 3.15 and 3.51 of [17]). Since  $K/K'$  is abelian (cf. 3.52 of [17]),  $G/K'$  is solvable and so  $K$  is perfect by the property of  $K$ . Note that  $K \subseteq H_G$ . Because  $G/H_G \cong (G/K)/(H_G/K)$  is solvable, it follows from part (iv) that  $H \in \mathcal{S}(G)$ .

(vi) As  $[G : H] < \infty$ ,  $G/H_G$  is finite and so there are finitely many elements  $x_1, \dots, x_n$  of  $G$  such that  $\langle x_1^*, \dots, x_n^* \rangle = G/H_G$  where  $x_i^* = x_i H_G$  for  $i = 1, \dots, n$ . If  $G$  is locally solvable, then finitely generated  $K = \langle x_1, \dots, x_n \rangle$  and the quotient  $G/H_G = KH_G/H_G \cong K/(K \cap H_G)$  are solvable, thus  $H \in \mathcal{S}(G)$  by part (iv).

Let  $[G : H]$  be square free. Then the factors of a composition series from  $H$  to  $G$  are simple groups with square-free orders. By Corollary 1 to Theorem 2.10 in Ch. 5 of [22], any group  $F \neq \{e\}$  of square-free order has a normal subgroup with the index being the least prime divisor of  $|F|$ . So the factors must have prime orders and hence  $H \in \mathcal{S}(G)$  by part (i).

Now suppose that  $[G : H]$  is odd or divisible by at most two distinct primes. Clearly so is  $|G/H_G|$  because we can view  $H_G$  as an intersection of finitely many conjugates of  $H$  and  $|G/H_G|$  must divide a power of  $[G : H]$  by Lemma 3.1(ii). Applying the well-known theorems of Feit–Thompson [3] and Burnside (cf. 8.5.3 of [16]), we then obtain the solvability of  $G/H_G$ . So  $H \in \mathcal{S}(G)$  by part (iv).

The proof of Theorem 3.1 is now complete.  $\square$

#### 4. THE MAIN RESULTS

**THEOREM 4.1.** *Let  $G$  be a group and (1.1) an exact  $m$ -cover of  $G$  by left cosets of subgroups  $G_1, \dots, G_k$ . Then, for any fixed  $i \in \{1, \dots, k\}$ ,*

$$k \geq m + f([G : G_i]) \quad \text{if } G_i \text{ contains some } H \in \mathcal{S}(G). \quad (4.1)$$

**PROOF.** Let  $1 \leq t \leq k$ ,  $H_t \in \mathcal{S}(G)$  and  $H_t \subseteq G_t$ . We use induction on  $[G : H_t]$  to show that  $k \geq m + f([G : G_t])$ .

If  $[G : H_t] = 1$ , then  $G_t = G$  and hence  $m + f([G : G_t]) = m = w_A(e) \leq k$ .

Now assume that  $[G : H_t] > 1$ . As  $H_t \in \mathcal{S}(G)$ , by Theorem 3.1(i) there exists a normal subgroup  $H$  of  $G$  for which  $H_t$  is subnormal in  $H$  and  $p = [G : H]$  is a prime. Write  $G/H = \{g_1H, \dots, g_pH\}$ . Put  $I_0 = \{1 \leq i \leq k : G_i \not\subseteq H\}$  and  $I_s = \{1 \leq i \leq k : a_i G_i \subseteq g_s H\}$  for  $s = 1, \dots, p$ . Clearly the union of these pairwise disjoint sets coincides with  $\{1, \dots, k\}$ . For any  $s = 1, \dots, p$ , by Lemma 2.1 we have  $a_i G_i \cap g_s H \neq \emptyset$  if and only if  $i \in I_0 \cup I_s$ , so  $I_0 \cup I_s \neq \emptyset$  and  $\{g_s^{-1} a_i G_i \cap H\}_{i \in I_0 \cup I_s}$  forms an exact  $m$ -cover of  $H$  by left cosets of subgroups  $G_i \cap H$  ( $i \in I_0 \cup I_s$ ).

Apparently  $H_t \subseteq G_t \cap H$  and  $[H : H_t] < [G : H_t]$ . Choose  $1 \leq s \leq p$  so that  $t \in I_0 \cup I_s$ . By the induction hypothesis,

$$|I_0 \cup I_s| \geq m + f([H : G_t \cap H]).$$

If  $t \in I_0$  then  $[G : G_t] = [G_t H : G_t] = [H : G_t \cap H]$  and hence

$$k \geq |I_0 \cup I_s| \geq m + f([H : G_t \cap H]) = m + f([G : G_t]).$$

In the case  $t \in I_s$  (whence  $G_t \subseteq H$ ), by Theorem 2.1 none of  $I_1, \dots, I_p$  is empty, thus

$$\begin{aligned} k &= |I_0| + |I_1| + \dots + |I_p| \geq |I_0 \cup I_s| + p - 1 \\ &\geq m + f([H : G_t]) + f([G : H]) = m + f([G : G_t]). \end{aligned}$$



This concludes the induction step.  $\square$

REMARK 4.1. Let (1.1) be an exact  $m$ -cover of a group  $G$ . Then  $[G : G_i]$  and  $[G : (G_i)_G]$  are finite. When  $G/(G_i)_G$  is solvable, we have  $H = (G_i)_G \in \mathcal{S}(G)$  by Theorem 3.1(iv), therefore  $k \geq m + f([G : G_i])$ .

COROLLARY 4.1. Let  $G$  be a group and (1.1) an exact  $m$ -cover of  $G$  by left cosets. Suppose that

$$\emptyset \neq I \subseteq \{1 \leq i \leq k : G_i \text{ contains a subgroup in } \mathcal{S}(G)\}.$$

Then for any subgroup  $K$  of  $G$  we have

$$(k-m)|I| \geq f\left(\left[K : K \cap \bigcap_{i \in I} G_i\right]\right), \quad (4.2)$$

thus

$$\left[G : \bigcap_{i \in I} G_i\right] \leq 2^{(k-m)|I|}. \quad (4.3)$$

PROOF. Let  $K$  be an arbitrary subgroup of  $G$  and  $t$  be any element of  $I$ . Put

$$J = \{1 \leq j \leq k : a_j G_j \cap a_t K \neq \emptyset\}.$$

Then  $t \in J$  and  $\{a_t^{-1} a_j G_j \cap K\}_{j \in J}$  forms an exact  $m$ -cover of  $K$  by left cosets of  $G_j \cap K$  ( $j \in J$ ). Let  $H_t$  be a subgroup in  $\mathcal{S}(G)$  contained in  $G_t$ . By Theorem 3.1(ii) we have  $H_t \cap K \in \mathcal{S}(K)$ . In view of Theorem 4.1,  $k \geq |J| \geq m + f([K : G_t \cap K])$ .

Write  $I = \{i_1, \dots, i_{|I|}\}$ . By the above,

$$\begin{aligned} k-m &\geq f([K : K \cap G_{i_1}]), \\ k-m &\geq f([K \cap G_{i_1} : K \cap G_{i_1} \cap G_{i_2}]), \\ &\dots\dots\dots, \\ k-m &\geq f([K \cap G_{i_1} \cap \dots \cap G_{i_{|I|-1}}, \\ &\quad K \cap G_{i_1} \cap \dots \cap G_{i_{|I|-1}} \cap G_{i_{|I|}}]). \end{aligned}$$

Adding these inequalities we then get the desired (4.2).

If we take  $K = G$ , then (4.2) gives that  $(k-m)|I| \geq f([G : \bigcap_{i \in I} G_i])$ , which implies (4.3) by (1.9). This completes the proof.  $\square$

COROLLARY 4.2. Let  $k \geq m > 0$  be integers. Then  $2^{k-m}$  is the maximal value that can be the index of a subgroup in a locally solvable group with an exact  $m$ -cover by  $k$  cosets one of which is a coset of the subgroup.

PROOF. Suppose that a locally solvable group  $G$  possesses an exact  $m$ -cover consisting of a coset  $C_1$  of subgroup  $G_1, \dots$ , a coset  $C_k$  of subgroup  $G_k$ . For  $i = 1, \dots, k$ , we let  $G_i^* = G_i$  if  $C_i$  is a left coset  $a_i G_i$ , and  $G_i^* = a_i^{-1} G_i a_i$  if  $C_i$  is a right coset  $G_i a_i$  of  $G_i$ . As  $\{a_i G_i^*\}_{i=1}^k$  forms an exact  $m$ -cover of  $G$ , each  $G_i^*$  has finite index in  $G$  and hence  $(G_i^*)_G \in \mathcal{S}(G)$  by Theorem 3.1(vi). In the light of Theorem 4.1, for each  $n_i = [G : G_i] = [G : G_i^*]$ , we have  $k \geq m + f(n_i)$  and hence  $n_i \leq 2^{k-m}$  by (1.9).

Now it suffices to notice that the following  $k$  residue classes

$$\underbrace{\mathbb{Z}, \dots, \mathbb{Z}}_{m-1}, 2^{k-m}\mathbb{Z}, 2^0 + 2\mathbb{Z}, \dots, 2^{k-m-1} + 2^{k-m}\mathbb{Z}$$

together form an exact  $m$ -cover of the infinite cyclic group  $\mathbb{Z}$  with the largest modulus being  $2^{k-m}$ . This follows from Example 1.2 in the case  $G = \mathbb{Z}$  and  $H = 2^{k-m}\mathbb{Z}$ .  $\square$

COROLLARY 4.3. Let (1.1) be an exact  $m$ -cover of a group  $G$  by cosets of subgroups  $G_1, \dots, G_k$ . Then  $k \geq m + f([G : G_i])$  if  $G_i$  contains a subnormal subgroup of  $G$  with index odd or square free or divisible by at most two distinct primes.

PROOF. Let  $H$  be any subnormal subgroup of  $G$  with  $[G : H]$  odd or square free or in the form  $p^\alpha q^\beta$  where  $p, q$  are primes and  $\alpha, \beta$  are non-negative integers. In view of Theorem 3.1(vi) we have  $H \in \mathcal{S}(G)$ . If  $G_i \supseteq H$ , then  $k \geq m + f([G : G_i])$  by Theorem 4.1. This concludes the proof.  $\square$

For an exact  $m$ -cover of a group  $G$ , if all the  $G_i$  are subnormal in  $G$  then we have a sharp lower bound of  $k$  in terms of the intersection  $H = \bigcap_{i=1}^k G_i$ .

THEOREM 4.2. Let  $G$  be a group and  $G_1, \dots, G_k$  be subnormal subgroups of  $G$  such that (1.1) forms an exact  $m$ -cover of  $G$  for some  $a_1, \dots, a_k \in G$ . Let  $K$  be any subgroup of  $G$  and set  $I(K) = \{1 \leq i \leq k : K \not\subseteq G_i\}$ .

(i) We have

$$k \geq m + d\left(K, K \cap \bigcap_{i=1}^k G_i\right). \quad (4.4)$$

(ii) If  $I(K) \neq \emptyset$ , then there exists an  $r \in I(K)$  and  $x_i \in K \setminus G_i$  for  $i \in I(K) \setminus \{r\}$ ,

such that

$$\left| \left\{ x_i \left( K \cap \bigcap_{s=1}^k G_s \right) : i \in I(K) \setminus \{r\} \right\} \right| \geq d\left(K, K \cap \bigcap_{s=1}^k G_s\right), \quad (4.5)$$

and hence

$$|\{1 \leq i \leq k : K \not\subseteq G_i\}| \geq 1 + d\left(K, K \cap \bigcap_{s=1}^k G_s\right). \quad (4.6)$$

PROOF. We use induction on the finite index  $[G : L]$  where  $L = \bigcap_{i=1}^k G_i$ .

If  $[G : L] = 1$ , then  $G_1 = \dots = G_k = G$ , so  $m + d(K, K \cap L) = m = w_{\mathcal{A}}(e) \leq k$  and  $I(K) = \emptyset$ .

Now let us proceed to the induction step and suppose that  $[G : L] > 1$ . Choose  $1 \leq j \leq k$  and a maximal normal proper subgroup  $H$  of  $G$  such that  $G_j$  is subnormal in  $H$ . Obviously  $\{1 \leq i \leq k : G_i \subseteq H\}$  can be partitioned into  $h = |G/H|$  sets

$$J(C) = \{1 \leq i \leq k : a_i G_i \subseteq C\} \quad (C \in G/H)$$

which are non-empty by Theorem 2.1. Let  $I = \{1 \leq i \leq k : G_i \not\subseteq H\}$ . If  $i \in I$ , then by Lemma 2.1 we have  $G_i H = G$  and hence  $a_i G_i \cap gH \neq \emptyset$  for all  $g \in G$ .

Let us take the first step. Set  $g_1 = a_j$  and  $I_1 = I \cup J(g_1 H)$ . Then  $j \in I_1$  and system  $\mathcal{A}_1 = \{g_1^{-1} a_i G_i \cap H\}_{i \in I_1}$  forms an exact  $m$ -cover of  $H_0 = H$  by left cosets of subnormal subgroups  $G_i \cap H_0$  ( $i \in I_1$ ) of  $H_0$ . Put

$$H_1 = H_0 \cap \bigcap_{i \in I_1} G_i = \bigcap_{i \in I_1} G_i \quad \text{and} \quad M_1 = \{i \in I_1 : K \cap H_0 \not\subseteq G_i \cap H_0\}.$$

Apparently  $M_1 \subseteq I(K)$ . Observe that

$$\left[ H_0 : \bigcap_{i \in I_1} (G_i \cap H_0) \right] = [H_0 : H_1] < [G : L].$$

By the induction hypothesis, we have

$$|I_1| \geq m + d\left(K \cap H_0, (K \cap H_0) \cap \bigcap_{i \in I_1} (G_i \cap H_0)\right) = m + d(K \cap H_0, K \cap H_1);$$

if  $M_1 \neq \emptyset$  (i.e.,  $K \cap H_0 \neq K \cap H_1$ ) then there is an  $r_1 \in M_1$  and  $x_i \in (K \cap H_0) \setminus (G_i \cap H_0) \subseteq K \setminus G_i$  for  $i \in M_1 \setminus \{r_1\}$  such that

$$|\{x_i(K \cap H_1) : i \in M_1 \setminus \{r_1\}\}| \geq d(K \cap H_0, K \cap H_1).$$

Suppose that we have found  $g_1, \dots, g_{s-1} \in G$  ( $s > 1$ ) and pairwise disjoint non-empty subsets  $I_1, \dots, I_{s-1}$  of  $\{1, \dots, k\}$  so that for each  $1 \leq t < s$ , either  $I_t \subseteq J(g_t H)$  or  $t = 1$ , and  $\mathcal{A}_t = \{g_t^{-1} a_i G_i \cap H_{t-1}\}_{i \in I_t}$  forms an exact  $m_t$ -cover of  $H_{t-1}$  by left cosets of subnormal subgroups  $G_i \cap H_{t-1}$  ( $i \in I_t$ ) of  $H_{t-1}$ , where  $0 < m_t \leq m_1 = m$  and we let

$$H_t = \bigcap_{i \in I_t} (G_i \cap H_{t-1}) = \bigcap_{i \in I_1 \cup \dots \cup I_t} G_i.$$

In the case  $I_s^* = \bigcup_{t=1}^{s-1} I_t \subset \{1, \dots, k\}$ , we proceed step  $s$  as follows. Select an element  $g_s$  in the union of those  $a_i G_i$  with  $i \notin I_s^*$ . Then  $m$  is greater than  $l_s = |\{i \in I_s^* : g_s \in a_i G_i\}|$ , and  $|\{i \in I_s^* : g_s x \in a_i G_i\}| = l_s$  for all  $x \in H_{s-1} = \bigcap_{i \in I_s^*} G_i$ . Put

$$I_s = \{1 \leq i \leq k : i \notin I_s^* \text{ \& } a_i G_i \cap g_s H_{s-1} \neq \emptyset\} \neq \emptyset.$$

For  $i \in I_s$  we have  $a_i G_i \cap g_s H \neq \emptyset$  and hence  $i \in J(g_s H)$  since  $I_s \cap I = \emptyset$ . Let  $m_s = m - l_s$ . Then  $\mathcal{A}_s = \{g_s^{-1} a_i G_i \cap H_{s-1}\}_{i \in I_s}$  forms an exact  $m_s$ -cover of  $H_{s-1}$  by left cosets of subnormal subgroups  $G_i \cap H_{s-1}$  ( $i \in I_s$ ) of  $H_{s-1}$ . Set

$$H_s = H_{s-1} \cap \bigcap_{i \in I_s} G_i = \bigcap_{i \in \bigcup_{t=1}^s I_t} G_i \quad \text{and} \quad M_s = \{i \in I_s : K \cap H_{s-1} \not\subseteq G_i \cap H_{s-1}\}.$$

Apparently  $M_s \subseteq I(K)$ , and  $M_s = \emptyset$  if and only if  $K \cap H_{s-1} = K \cap H_s$ . In the light of the induction hypothesis,

$$|I_s| \geq m_s + d\left(K \cap H_{s-1}, K \cap H_{s-1} \cap \bigcap_{i \in I_s} (G_i \cap H_{s-1})\right) \geq 1 + d(K \cap H_{s-1}, K \cap H_s);$$

if  $M_s \neq \emptyset$  then there is an  $r_s \in M_s$  and  $x_i \in (K \cap H_{s-1}) \setminus (G_i \cap H_{s-1}) \subseteq K \setminus G_i$  for  $i \in M_s \setminus \{r_s\}$  such that

$$|\{x_i(K \cap H_s) : i \in M_s \setminus \{r_s\}\}| \geq d(K \cap H_{s-1}, K \cap H_s).$$

Since  $h = |G/H| > 1$  we have  $I_1 \subset \{1, \dots, k\}$ . As  $\{1, \dots, k\}$  is a finite set the above process will terminate after  $n$  steps where  $1 < n \leq k$ . Thus  $\bigcup_{s=1}^n I_s = \{1, \dots, k\}$  and  $H_n = \bigcap_{i=1}^k G_i = L$ . Because

$$\bigcup_{C \in G/H} J(C) = \{1 \leq i \leq k : G_i \subseteq H\} = \bigcup_{s=1}^n \{i \in I_s : G_i \subseteq H\} \subseteq \bigcup_{s=1}^n J(g_s H),$$

we have  $n \geq h = |G/H| \geq l = [K : H \cap K] \geq 1 + d(K, H \cap K)$ . Therefore

$$\begin{aligned} k &= |I_1| + \sum_{s=2}^n |I_s| \geq m + d(K \cap H_0, K \cap H_1) + \sum_{s=2}^n (1 + d(K \cap H_{s-1}, K \cap H_s)) \\ &\geq m + d(K, H \cap K) + d(K \cap H_0, K \cap H_n) = m + d(K, K \cap L). \end{aligned}$$

Let

$$M = \bigcup_{\substack{1 \leq s \leq n \\ K \cap H_{s-1} \neq K \cap H_s}} (M_s \setminus \{r_s\}).$$

By the above  $M \subseteq I(K)$ . For  $x, y \in K$ , if  $1 \leq s \leq n$  and  $x(K \cap H_s) \neq y(K \cap H_s)$ , then  $x(K \cap L) \neq y(K \cap L)$  (otherwise  $x^{-1}y \in K \cap L \subseteq K \cap H_s$ ). If  $1 \leq t < s \leq n$ ,  $x \in K \cap H_{s-1}$  and  $y \in (K \cap H_{t-1}) \setminus (G_i \cap H_{t-1})$  for some  $i \in I_t$ , then  $x(K \cap L) \subseteq H_{s-1} \subseteq H_t \subseteq G_i \cap H_{t-1}$  and  $y(K \cap L) \cap (G_i \cap H_{t-1}) = \emptyset$ . Therefore

$$\begin{aligned} |\{x_i(K \cap L) : i \in M\}| &= \sum_{\substack{1 \leq s \leq n \\ K \cap H_{s-1} \neq K \cap H_s}} |\{x_i(K \cap L) : i \in M_s \setminus \{r_s\}\}| \\ &\geq \sum_{\substack{1 \leq s \leq n \\ K \cap H_{s-1} \neq K \cap H_s}} |\{x_i(K \cap H_s) : i \in M_s \setminus \{r_s\}\}| \\ &\geq \sum_{\substack{1 \leq s \leq n \\ K \cap H_{s-1} \neq K \cap H_s}} d(K \cap H_{s-1}, K \cap H_s) = \sum_{s=1}^n d(K \cap H_{s-1}, K \cap H_s) \\ &= d(K \cap H_0, K \cap H_n) = d(H \cap K, K \cap L). \end{aligned}$$

If  $1 \leq s \leq n$  and  $M_s = \emptyset$ , then we let  $r_s$  be an element of  $I_s$ . Clearly  $M' = \{r_s : 1 < s \leq l\}$  has cardinality  $l-1$ , also  $M' \cap M = \emptyset$  and  $M \cup M' \subseteq I(K) \setminus \{r_1\}$ . If  $1 < s \leq l$ , then  $G_{r_s} \subseteq H$  since  $r_s \in I_s \subseteq J(g_s H)$ . Write  $K/(H \cap K) = \{H \cap K, b_1(H \cap K), \dots, b_l(H \cap K)\}$ . When  $1 < s \leq l$ , we have  $b_s \notin H$  (otherwise  $b_s \in H \cap K$ , which is impossible) and hence  $x_{r_s} = b_s \in K \setminus G_{r_s}$ . Thus  $M \cup M' \subseteq I(K) \setminus \{r_1\}$ . Recall that  $x_i \in K \cap H_{s-1} \subseteq H \cap K$  for  $i \in M_s \setminus \{r_s\}$ . For  $x, y \in K$  with  $x(H \cap K) \neq y(H \cap K)$ , obviously  $x(K \cap L) \neq y(K \cap L)$ . For  $i \in (I(K) \setminus \{r_1\}) \setminus (M \cup M')$  let  $x_i$  be any element of  $K \setminus G_i$ . Then

$$\begin{aligned} |\{x_i(K \cap L) : i \in I(K) \setminus \{r_1\}\}| \\ &\geq |\{x_i(K \cap L) : i \in M \cup M'\}| = |\{x_i(K \cap L) : i \in M'\}| + |\{x_i(K \cap L) : i \in M\}| \\ &\geq l-1 + d(H \cap K, K \cap L) \geq d(K, H \cap K) + d(H \cap K, K \cap L) = d(K, K \cap L) \end{aligned}$$

and therefore  $|I(K)| \geq 1 + d(K, K \cap L)$ .

The proof by induction is now complete.  $\square$

REMARK 4.2. (a) Clearly  $I(K) \subseteq \{1 \leq i \leq k : G_i \neq G\}$ , so we cannot substitute  $m$  ( $\geq |\{1 \leq i \leq k : G_i = G\}|$ ) for the first term 1 on the right-hand side of (4.6). (b) Let  $H$  be a subnormal subgroup of finite index in group  $G$ . If (1.1) forms an exact  $m$ -cover of  $G$  with  $G_1, \dots, G_k$  subnormal in  $G$  and  $\bigcap_{i=1}^k G_i = H$ , then by taking  $K = G$  in Theorem 4.2(i) we get the inequality  $k \geq m + d(G, H)$ . On the other hand, by Example 1.2 there indeed exists an exact  $m$ -cover (1.1) of  $G$  with all the  $G_i$  subnormal in  $G$ ,  $\bigcap_{i=1}^k G_i = H$  and  $k = m + d(G, H)$ .

COROLLARY 4.4. Let  $G$  be a group and  $H, K$  be subnormal subgroups of  $G$  with finite index. Let  $G_1, \dots, G_k$  be subnormal subgroups of  $G$  for which all the  $G_i$  contain  $H$  but  $I(K) = \{1 \leq i \leq k : K \not\subseteq G_i\} \neq \emptyset$ . Suppose that  $X = \bigcup_{i=1}^k a_i G_i$  is a union of left cosets of  $K$  and  $w_A(x) = m$  for all  $x \in X$ , where  $a_1, \dots, a_k \in G$ . Then there are  $C_i \in K/(H \cap K)$  for those  $i \in I(K)$  such that  $C_i \cap G_i \neq \emptyset$  for a unique  $i \in I(K)$  and that

$$|\{C_i : i \in I(K)\}| \geq 1 + d\left(K, K \cap \bigcap_{i=1}^k G_i\right). \quad (4.7)$$

PROOF. The set  $G \setminus X$  can be written as a union of finitely many distinct left cosets  $b_1K, \dots, b_lK$  of  $K$ . Let  $G_{k+1} = \dots = G_{k+lm} = K$  and  $a_{k+(j-1)m+1} = \dots = a_{k+jm} = b_j$  for  $j = 1, \dots, l$ . Clearly  $\mathcal{A}' = \{a_i G_i\}_{i=1}^{k+lm}$  forms an exact  $m$ -cover of  $G$  by left cosets of subnormal subgroups of  $G$ . Observe that  $\bigcap_{i=1}^{k+lm} G_i = K \cap L$ , where  $L = \bigcap_{i=1}^k G_i$ . Also,  $\{1 \leq i \leq k+lm : K \not\subseteq G_i\} = I(K) \neq \emptyset$ . In the light of Theorem 4.2(ii), there is an  $r \in I(K)$  and  $x_i \in K \setminus G_i$  for  $i \in I(K) \setminus \{r\}$  such that  $|\{x_i(K \cap L) : i \in I(K) \setminus \{r\}\}| \geq d(K, K \cap L)$ . Put  $C_r = H \cap K$  and  $C_i = x_i(H \cap K)$  for  $i \in I(K) \setminus \{r\}$ . Note that  $e \in C_r \cap G_r$ . For  $i \in I(K) \setminus \{r\}$  we have  $C_i \cap G_i = \emptyset$  because  $C_i \subseteq x_i G_i$  and  $x_i \notin G_i$ . If  $s$  and  $t$  are distinct elements of  $I(K) \setminus \{r\}$ , then

$$C_s = C_t \Rightarrow x_s^{-1}x_t \in H \cap K \subseteq K \cap L \Rightarrow x_s(K \cap L) = x_t(K \cap L).$$

So

$$\begin{aligned} |\{C_i : i \in I(K)\}| &= 1 + |\{C_i : i \in I(K) \setminus \{r\}\}| \\ &\geq 1 + |\{x_i(K \cap L) : i \in I(K) \setminus \{r\}\}| \geq 1 + d(K, K \cap L). \end{aligned}$$

This completes the proof.  $\square$

REMARK 4.3. When  $m = 1$  and  $G_1, \dots, G_k, K$  are normal in  $G$ , the inequality  $k \geq 1 + d(K, K \cap \bigcap_{i=1}^k G_i)$  was obtained by Korec [7] in another way. Theorem 9 of [18] is essentially Corollary 4.4 in the case  $m = 1$  and  $X = G$ .

COROLLARY 4.5. *Let  $m$  be a positive integer and  $H$  a subgroup of a group  $G$  with  $[G : H]$  finite.*

- (i) *If  $G$  is locally nilpotent, then  $m + f([G : H])$  is the least positive integer  $k$  such that there exists an exact  $m$ -cover of  $G$  by  $k$  left cosets of subgroups whose intersection is  $H$ .*
- (ii) *Providing  $H$  is subnormal in  $G$ , if  $G$  is locally solvable, or  $[G : H]$  odd or square free or in the form  $p^\alpha q^\beta$  where  $p, q$  are distinct primes and  $\alpha, \beta$  are non-negative integers, then  $m + f([G : H])$  is the smallest  $k \in \mathbb{Z}^+$  such that there exists an exact  $m$ -cover (1.1) of  $G$  with all the  $G_i$  subnormal in  $G$  and  $\bigcap_{i=1}^k G_i$  equal to  $H$ .*

PROOF. In view of Remark 4.2(b), it suffices to make the following observations:

- (i) When  $G$  is locally nilpotent, by Theorem 7 of [18]  $H$  is in  $\mathcal{S}(G)$  and any subgroup containing  $H$  is subnormal in  $G$ .
- (ii) Suppose that  $H$  is subnormal in  $G$ , and that  $G$  is locally solvable or  $[G : H]$  is odd or square free or divisible by at most two distinct primes. Then  $d(G, H) = f([G : H])$  by Theorem 3.1(vi).  $\square$

COROLLARY 4.6. *Let  $H$  be a subnormal subgroup of a group  $G$  with  $[G : H] < \infty$ . Then  $H$  is normal in  $G$  if and only if*

$$|N_G(H)/H| + d(H, H_G) \geq [G : H] \quad (4.8)$$

where  $N_G(H)$  denotes the normalizer of  $H$  in  $G$ .

PROOF. If  $H$  is normal in  $G$  then (4.8) holds because  $N_G(H) = G$  and  $H_G = H$ .

Below we assume that  $H$  is not normal in  $G$ . Let  $k = [G : H]$ , and  $\{Ha_1, \dots, Ha_k\}$  be a partition of  $G$  into right cosets of  $H$ . Clearly  $G_i = a_i^{-1}Ha_i$  is subnormal in  $G$  for each

$i = 1, \dots, k$ . As mentioned in Example 1.1 system (1.1) forms a disjoint cover of  $G$  with  $\bigcap_{i=1}^k G_i = H_G$ . Since  $H \supset H_G = \bigcap_{i=1}^k G_i$ , it follows from Theorem 4.2(ii) that

$$|\{1 \leq i \leq k : H \not\subseteq G_i\}| \geq 1 + d\left(H, H \cap \bigcap_{i=1}^k G_i\right) = 1 + d(H, H_G).$$

For  $i = 1, \dots, k$ , as  $[G : G_i] = [G : H] < \infty$ ,  $H \subseteq G_i$  if and only if  $H = G_i$ . We have  $|N_G(H) : H| = |\{1 \leq i \leq k : G_i = H\}|$ , because

$$\begin{aligned} N_G(H) &= \{g \in G : g^{-1}Hg = H\} = \bigcup_{i=1}^k \{ha_i : h \in H \text{ \& } (ha_i)^{-1}Hha_i = H\} \\ &= \bigcup_{i=1}^k \{ha_i : h \in H \text{ \& } a_i^{-1}Ha_i = H\} = \bigcup_{\substack{i=1 \\ a_i^{-1}Ha_i=H}}^k Ha_i. \end{aligned}$$

Therefore

$$\begin{aligned} d(H, H_G) &< |\{1 \leq i \leq k : H \not\subseteq G_i\}| = |\{1 \leq i \leq k : G_i \neq H\}| \\ &= k - |\{1 \leq i \leq k : G_i = H\}| = [G : H] - |N_G(H)/H|. \end{aligned}$$

Consequently (4.8) fails to hold. We are done.  $\square$

REMARK 4.4. Let  $G$  be a group and  $H$  be a subnormal subgroup of  $G$  with finite index. By Corollary 3 of [18],  $[G : H] \geq 1 + d(G, H_G)$  and consequently  $|G/H_G| \leq 2^{[G:H]-1}$ . In view of Corollary 4.6 and (1.9), if  $H$  is not normal in  $G$  then  $|H/H_G| \leq 2^{d(H, H_G)} \leq 2^{[G:H]-1-|N_G(H)/H|}$ . It seems that  $|N_G(H)/H| \geq d(G, H)$ .

#### ACKNOWLEDGEMENTS

This research is supported by the Teaching and Research Award Fund for Outstanding Young Teachers in Higher Education Institutions of MOE, and the National Natural Science Foundation of the People's Republic of China.

#### REFERENCES

1. M. A. Berger, A. Felzenbaum and A. S. Fraenkel, Mycielski–Sierpiński conjecture and Korec–Znám result, *Colloq. Math.*, **56** (1988), 241–249, MR 90d:20044.
2. P. Erdős, On integers of the form  $2^k + p$  and some related problems, *Summa Brasil. Math.*, **2** (1950), 113–123, MR 13, 437.
3. W. Feit and J. G. Thompson, Solvability of groups of odd order, *Pac. J. Math.*, **13** (1963), 775–1029, MR 29#3538.
4. R. K. Guy, *Unsolved Problems in Number Theory*, 2nd Ed, Springer-Verlag, New York, 1994, Sections A19, B21, E23, F13, F14.
5. M. Hejrány and Š. Znám, Coset decomposition of Abelian groups, *Acta Fac. Rerum Nat. Univ. Comen. Math. Publ.*, **25** (1971), 15–19, MR 46#3644.
6. I. Korec, On a generalization of Mycielski's and Znám's conjectures about coset decomposition of Abelian groups, *Fundam. Math.*, **85** (1974), 41–47, MR 50#10025.
7. I. Korec, On number of cosets in nonnatural disjoint covering systems, in: *Number Theory*, Vol. I, Colloq. Math. Soc. János Bolyai, North-Holland, Amsterdam, 1990, **51**, pp. 265–278, MR 91k:11016.

8. I. Korec and Š. Znáám, On disjoint covering of groups by their cosets, *Math. Slovaca*, **27** (1977), 3–7, MR 58#5260.
9. J. Mycielski and W. Sierpiński, Sur une propriété des ensembles linéaires, *Fundam. Math.*, **58** (1966), 143–147, MR 34#4140.
10. B. H. Neumann, Groups covered by permutable subsets, *J. London Math. Soc.*, **29** (1954), 236–248, MR 15, 931.
11. B. H. Neumann, Groups covered by finitely many cosets, *Publ. Math. Debrecen*, **3** (1954), 227–242, MR 17, 234.
12. B. Novák and Š. Znáám, Disjoint covering systems, *Am. Math. Monthly*, **81** (1974), 42–45, MR 48#10958.
13. M. M. Parmenter, Exact covering systems for groups, *Fundam. Math.*, **123** (1984), 133–136, MR 86h:20029.
14. Š. Porubský, On  $m$  times covering systems of congruences, *Acta Arith.*, **29** (1976), 159–169, MR 53#2884.
15. Š. Porubský, *Results and Problems on Covering Systems of Residue Classes*, Mitt. Math. Sem. Giessen, Heft 150, Giessen Univ., 1981, pp. 1–85, MR 83b:10068.
16. D. J. S. Robinson, *A Course in the Theory of Groups*, Springer-Verlag, New York, 1982.
17. J. S. Rose, *A Course on Group Theory*, Cambridge University Press, Cambridge, U.K., 1978.
18. Z. W. Sun, Finite coverings of groups, *Fundam. Math.*, **134** (1990), 37–53, MR 91g:20031.
19. Z. W. Sun, On exactly  $m$  times covers, *Isr. J. Math.*, **77** (1992), 345–348, MR 93k:11007.
20. Z. W. Sun, Covering the integers by arithmetic sequences II, *Trans. Am. Math. Soc.*, **348** (1996), 4279–4320, MR 97c:11011.
21. Z. W. Sun, Exact  $m$ -covers and the linear form  $\sum_{s=1}^k x_s/n_s$ , *Acta Arith.*, **81** (1997), 175–198, MR 98h:11019.
22. M. Suzuki, *Group Theory I–II*, (Grundlehren der mathematischen Wissenschaften; 247–248), Springer-Verlag, New York, 1982, 1986.
23. M. J. Tomkinson, Groups covered by finitely many cosets or subgroups, *Commun. Algebra*, **15** (1987), 845–859, MR 88c:20049.
24. M. Z. Zhang, On irreducible exactly  $m$  times covering system of residue classes, *J. Sichuan Univ. (Nat. Sci. Ed.)*, **28** (1991), 403–408, MR 92j:11001.
25. Š. Znáám, On Mycielski’s problem on systems of arithmetic progressions, *Colloq. Math.*, **15** (1966), 201–204, MR 34#134.
26. Š. Znáám, On exactly covering systems of arithmetic sequences, in: *Number Theory (Colloq. János Bolyai Math. Soc., Debrecen 1968)*, North-Holland, Amsterdam, 1970, pp. 221–225, MR 42#7586.
27. Š. Znáám, A survey of covering systems of congruences, *Acta Math. Univ. Comenian.*, **40/41** (1982), 59–79, MR 84e:10004.

Received 10 January 2000 and accepted 3 November 2000

ZHI-WEI SUN  
 Department of Mathematics,  
 Nanjing University,  
 Nanjing 210093,  
 The People’s Republic of China  
 E-mail: zwsun@nju.edu.cn